arXiv:1207.0756v1 [nucl-th] 3 Jul 2012

Description of He isotopes using the particle-particle random-phase approximation model

B. Laurent[1, *] and N. Vinh Mau[2, 1]
[1] *CEA, DAM, DIF F-91297 Arpajon Cedex, France*
[2] *Institut de Physique Nucléaire, IN2P3-CNRS, Université Paris-Sud, F-91406 Orsay Cedex, France*

(Dated: July 4, 2012)

A particle-particle random phase approximation (RPA) model was applied to study the properties of the ^{6,8,9,10}He isotopes. The RPA model applied takes into account the two-body correlations in the ^{8}He core. Therefore, it gives a better description of energies and amplitudes than models that assume a closed neutron shell (or subshell) and provides the opportunity to make a simultaneous study of the ^{6}He, ^{8}He, ^{9}He and ^{10}He nuclei. Both ground-state and excited-states calculated properties are discussed and compared to experimental data. These studies suggest the same $2s$-$1p_{1/2}$ shell inversion in ^{9}He as in ^{11}Be, ^{10}Li and ^{13}Be, and propose a contribution of the $(1p_{3/2})^{-2} - (1p_{1/2})^2$ and the $(1p_{3/2})^{-2} - (2s)^2$ configurations in the ^{8}He core.

PACS numbers: 21.10.Re, 21.60.Jz, 27.20.+n

I. INTRODUCTION

Despite their small number of nucleons, the light nuclei of the nuclide chart provide a unique opportunity to investigate complex phenomena such as halo nuclei, cluster configurations, shell inversions, and give access to nuclear systems beyond the neutron dripline [1]. This region thus challenges the theory in its capacity to reproduce the wide variety of these phenomena. In that sense, the unbound helium isotopes, ^{9}He and ^{10}He have been extensively studied both experimentally [2–11] and theoretically [5, 12–16]. The first experiments on ^{9}He showed a $1/2^-$ state in the range of 1-2 MeV interpreted as its ground state [2–4, 17]. This property of a $1/2^-$ ground state was also the result of most of the calculations [18, 19]. A latter double proton knockout experiment however shows the existence of a lower state [5], close to the n+^{8}He threshold, compatible with a possible $1/2^+$ state. This assignment has been comforted by a shell model calculation [5]. Yet, recent experimental studies [9, 10] do not confirm such a low state.

Such a situation of a $1/2^+$ ground state with a $1/2^-$ excited state reveals an inversion of the $2s$ and $1p_{1/2}$ shells when ^{9}He is described as a neutron added to a core of ^{8}He. Such an inversion is well known in ^{11}Be [20], ^{10}Li [21] and ^{13}Be [22]. In ^{10}Li and ^{13}Be this inversion was first suggested theoretically in order to reproduce the two-neutron separation energy in ^{11}Li and ^{14}Be respectively [21, 23, 24], and then experimentally confirmed [7, 22]. These two-neutron separation energies were calculated in a two-neutron RPA model where ^{11}Li and ^{14}Be were considered as two neutrons added to an inert core of ^{9}Li and ^{12}Be respectively. In the present paper, the same two-neutron RPA model depicting ^{10}He as two neutrons added to an inert core of ^{8}He is used. This model gives the same system of equations for ^{10}He and ^{6}He; the latter being described as two-neutrons subtracted from a ^{8}He core. Finally these calculations provide, among other properties, the two-neutron separation energies (S_{2n}) of ^{8}He and ^{10}He, the 0^+ states in ^{10}He and the wave functions of ^{6}He and ^{10}He. The analysis of these wave functions or more exactly of the two-neutron RPA amplitudes include information about the core wave function.

In section II the two-neutron RPA model and the inputs used in the calculation are briefly presented. The results are given in section III and the conclusions are compiled in the final section.

II. THE MODEL

Only a brief summary of the used two-neutron RPA model is given. More details can be found in reference [25]. A core of ^{8}He is assumed which, in the Hartree-Fock model, is a closed shell nucleus with the last neutrons filling the $1p_{3/2}$ shell. Two neutrons are then added to, or subtracted from this core to describe ^{10}He and ^{6}He as eigenstates of a single system of equations. Indeed eigenvalues and eigenvectors of the RPA matrix yield the energies of ^{6}He and ^{10}He referred to the core energy and the two-body amplitudes in ^{6}He and ^{10}He with the following definition for a solution of rank n:

$$X^{(n)}(a) = \langle \psi_n(^{10}He) | A_a^+ | \psi_0(^{8}He) \rangle, \tag{1}$$

$$Y^{(n)}(a) = \langle \psi_n(^{6}He) | A_a | \psi_0(^{8}He) \rangle, \tag{2}$$

with the orthonormalization relations:

$$\sum_{\substack{occupied\\states}} |X(a)|^2 - \sum_{\substack{unoccupied\\states}} |X_(a)|^2 = 1,$$

$$\sum_{\substack{unoccupied\\states}} |Y(a)|^2 - \sum_{\substack{occupied\\states}} |Y(a)|^2 = 1, \tag{3}$$

---

[*] Corresponding author: benoit.laurent@cea.fr

where A_a^+ and A_a are respectively the creation and annihilation operators of two neutrons in state $\boldsymbol{a}$ formed of two occupied or unoccupied states in the Hartree-Fock core. ψ_0 is the correlated wave function of the ^{8}He core. Then, for example, a non zero amplitude $\boldsymbol{X}$ for $\boldsymbol{a}$, representing two occupied states, means that in the core wave function there is a configuration with two holes in that state. Similarly, if $\boldsymbol{Y}$ is non zero for a state $\boldsymbol{a}$ formed of two unoccupied states, then the ^{8}He wave function contains a configuration with two particles in that state $\boldsymbol{a}$.

Furthermore the eigenvalues give the energies of ^{6}He and ^{8}He referred to the core energy so that the two neutron separation energy in ^{8}He and ^{10}He are quite straightforward to reach. To achieve this calculation the D1S Gogny interaction [26, 27] is used as an effective nucleon-nucleon interaction. Also, the single neutron energy in the core of ^{8}He is needed. For these calculations, a Saxon-Woods potential is chosen for the neutron-core potential, with an additional surface term which takes into account the neutron-phonon couplings. Then the one-body potential is written as:

$$V_{nc}(r) = -V_{NZ}\left(f(r) - 0.44r_0^2(\mathbf{l.s})\frac{1}{r}\frac{df(r)}{dr}\right) + 16a^2\alpha_n\left(\frac{df(r)}{dr}\right)^2, \quad (4)$$

$$with\ f(r) = \left(1 + exp\left[\frac{r-R_0}{a}\right]\right)^{-1}. \quad (5)$$

The following parameters given by Bohr and Mottelson [28], are used:

$$V_{NZ} = U_0 - U_\tau\frac{N-Z}{A_c} = \left(51 - 33\frac{N-Z}{A_c}\right)\ MeV, \quad (6)$$

$$R_0 = r_0A_c^{1/3},\ r_0 = 1.27\ fm, \quad (7)$$

where N, Z, A_c refer to the core nucleus. The diffusivity is set to a larger value, $a = 0.75$ fm, to account for the diffuse surface of such a light nucleus. The parameter α_n is fitted according to the one-neutron separation energy in ^{8}He for the $1p_{3/2}$ state. For the $2s$ and $1p_{1/2}$ states, it is varied in order to reproduce the considered energies. For higher states, the parameter is set to zero since it is known that the neutron-phonon couplings effects on the potential are very small for states far from the Fermi surface.

III. RESULTS

All experiments agree on the presence in the ^{9}He spectrum of a $1/2^-$ state in the vicinity of 1-1.27 MeV energy range. This state was, up to L. Chen and collaborators' experiment [5], considered as the ground state of ^{9}He. This was attested by several shell model calculations. However L. Chen et *al.* experiment and most of the recent ones agree that the $1/2^-$ state is an excited state and that the ground state is an unbound virtual $1/2^+$ state, with an energy very close to the $n+^8$He threshold. On the other end, there is an ambiguity in the interpretation of experiments on ^{10}He. Indeed until recently the resonance found at about 1 MeV above the $2n+^8$He threshold was considered to be the ground state of ^{10}He. However there is a recent suggestion by [19] that this resonance corresponds to a 0^+ excited state and that the ground state is close to the threshold. The use of the two-neutron RPA model assuming a core of ^{8}He gives the opportunity to make a simultaneous study of ^{6}He, ^{8}He, ^{9}He and ^{10}He where the choice of parameters defining ^{9}He states determines the states of the other three nuclei without any further fitting possibility. The calculations proceed as follows: the $1p_{3/2}$ energy is set to the experimental energy of 2.57 MeV. The $1p_{1/2}$ energy is varied in a domain compatible with measurements. For each adopted value a series of calculations varying the $2s$ shell energy is made. The results are presented for $\epsilon(1p_{1/2}) = 1.25$ MeV, which is the value given in most of the experiments' analysis and for $\epsilon(1p_{1/2}) = 0.92$ MeV, which is not excluded by experiments. These results are given in tables I and II where the following observables are calculated: $S_{2n}(^8He)$, $S_{2n}(^{10}He)$ and $E(0_2^+)$. The latter corresponds to the energy of the first excited 0^+ state in ^{10}He relative to the $n + n+^8$He threshold. For $\epsilon(1p_{1/2}) = 1.25$ MeV and $\epsilon(2s) \approx 0.4 - 0.3$ MeV, ^{10}He is unbound with a small value of S_{2n} in agreement with the assumption that the resonance at about 1 MeV is not the ground state but a 0^+ excited state. Yet the energy of this excited state is too high. Furthermore $S_{2n}(^8He)$ is slightly too large compared to the experimental measured value of 2.14 MeV [29]. If we assume the $2s$ shell to be higher than the $1p_{1/2}$ one, for any value of $\epsilon(p_{1/2})$, the results are in complete disagreement with measurements.

TABLE I. Results in terms of $\epsilon(2s)$ when $\epsilon(1p_{1/2}) = 1.25$ MeV

$\epsilon(2s)$	0.78	0.45	0.29	0.192
$S_{2n}(^8He)$	2.52	2.48	2.48	2.43
$S_{2n}(^{10}He)$	−1.33	−0.53	−0.11	0.12
$E(0_2^+)$	2.3	2.3	2.3	2.3

TABLE II. Results in terms of $\epsilon(2s)$ when $\epsilon(1p_{1/2}) = 0.92$ MeV

$\epsilon(2s)$	0.785	0.45	0.29	0.19
$S_{2n}(^8He)$	2.36	2.31	2.29	2.24
$S_{2n}(^{10}He)$	−1.11	−0.44	−0.07	0.22
$E(0_2^+)$	1.6	1.4	1.4	1.4

On the other hand, in table II for $\epsilon(1p_{1/2}) = 0.92$ MeV a general agreement is observed for all calculated quantities though the most accurate are obtained for $\epsilon(2s) = 0.2 - 0.3$ MeV. This value of 0.92 MeV is slightly smaller than that commonly used in the interpretation of experiments. However if one looks, for example, at the figure



FIG. 1. Relative energy spectrum for ^{8}He+n system in C(^{14}B,^{8}He+n)X reaction. The dotted line represents the uncorrelated background. The thin solid line is the virtual s-state with $a_s = 0$ fm, while the dashed line is the p-resonance. The thicker solid line is the overall adjustment compared to the data (dots). The insert presents the best adjustment as a function of the contribution of each p-resonance and s-state. (Extracted from [9, 30])

1 given as fig. 1 in ref. [30], a $p_{1/2}$ resonance at about 0.9 MeV would give the same or even better agreement as 1.27 MeV. With these energies of $2s$ and $1p_{1/2}$ states, the two-neutron separation energy in ^{8}He is very close to the measured value of 2.14 MeV while the energy of the excited 0^+ state in ^{10}He is close to the energy of 1 MeV deduced from experiments. The advantage of this model is that, once the single neutron energies are fixed, the properties of all nuclei are given, without any possibility to introduce further parameters to modify the results for one or the other of the nuclei.

The RPA equations also yield the wave functions of ^{6}He and ^{10}He through the two-neutron amplitudes defined in Eqs. (1-3). In the best case where $\epsilon(1p_{1/2}) =$ 0.92 MeV, $\epsilon(2s) = 0.3$ MeV, the following amplitudes for ^{10}He ground state are obtained:

$$X(2s^2) = 0.98,$$
$$X(p^2_{1/2}) = -0.28,$$
$$X(p^2_{3/2}) = -0.28.$$

The main contribution comes from the $(2s)^2$ configuration for the two last neutrons but the $(1p_{3/2})^2$ shows a significant amplitude which means that in the core of ^{8}He there is a non negligible contribution of a 2 holes-2 particles configuration with the holes in the $p_{3/2}$-shell.

For ^{6}He, the obtained amplitudes are:

$$Y(p^{-2}_{3/2}) = -1.08,$$
$$Y(p^{-2}_{1/2}) = -0.21,$$
$$Y(2s^{-2}) = 0.20.$$

These values of the Y amplitudes indicate again a contribution of a 2 holes-2 particles configuration with the two particles in the $1p_{1/2}$ or $2s$ shells. Then from these two results it can be understood that ^{8}He is not a pure closed shell nucleus but has a contribution of $(1p_{3/2})^{-2} - (1p_{1/2})^2$ or $(1p_{3/2})^{-2} - (2s)^2$ configurations. The 0^+ excited state of ^{10}He appears to have mostly a $(1p_{1/2})^2$ configuration with some mixture of $(2s)^2$, with the respective amplitudes of 0.96 and 0.25.

IV. CONCLUSIONS

In a two-neutron RPA model we have made a simultaneous study of the ^{6}He, ^{8}He, ^{9}He, ^{10}He isotopes. The results show that, to reach the best two-neutron separation energy in ^{8}He, a good interpretation of experiments for ^{10}He, ^{9}He should have a $1/2^+$ unbound ground state close to the threshold and a $1/2^-$ excited state close to the measured resonance often assumed as the ground state. The corresponding unbound ground state of ^{10}He is close to the threshold and the observed resonance at 1 MeV is an excited 0^+ state, not the ground state as usually assumed. Furthermore the RPA amplitudes obtained for ^{6}He and ^{10}He show that the core of ^{8}He is not a pure closed $(1p_{3/2})$ shell nucleus but has a qualitatively important mixture of $(1p_{3/2})^{-2} - (1p_{1/2})^2$ and $(1p_{3/2})^{-2} - (2s)^2$ configurations. The wave functions of ^{6}He and ^{10}He show also some mixtures of various configurations with still a strong component on two holes in the $p_{3/2}$-shell for ^{6}He and on two particles in the $2s$-shell or $p_{1/2}$-shell for respectively the ground state and the 0^+_2 excited state in ^{10}He.

---

[1] B. Jonson, Phys. Rep. **389**, 1 (2004).

[2] K. K. Seth, M. Artuso, D. Barlow, S. Iversen, M. Kaletka, H. Nann, B. Parker, and R. Soundranayagam, Phys. Rev. Lett. **58**, 1930 (1987).

[3] H. G. Bohlen, B. Gebauer, D. Kolbert, W. von Oertzen, E. Stiliaris, M. Wilpert, and T. Wilpert, Z. Phys. A **330**, 227 (1988).

[4] W. von Oertzen, H. Bohlen, B. Gebauer, M. von Lucke-Petsch, A. Ostrowski, C. Seyfert, T. Stolla,

M. Wilpert, T. Wilpert, D. Alexandrov, A. Korsheninnikov, I. Mukha, A. Ogloblin, R. Kalpakchieva, Y. Penionzhkevich, S. Piskor, S. Grimes, and T. Massey, Nucl. Phys. A **588**, c129 (1995).

[5] L. Chen and et *al.* , Phys. Lett. B **505**, 21 (2001).

[6] G. V. Rogachev, V. Z. Goldberg, J. J. Kolata, G. Chubarian, D. Aleksandrov, A. Fomichev, M. S. Golovkov, Y. T. Oganessian, A. Rodin, B. Skorodumov, R. S. Slepnev, G. Ter-Akopian, W. H. Trzaska, and R. Wolski, Phys. Rev. C **67**, 041603 (2003).

[7] S. Fortier, E. Tryggestad, E. Rich, D. Beaumel, E. Becheva, Y. Blumenfeld, F. Delaunay, A. Drouart, A. Fomichev, N. Frascaria, S. Gales, L. Gaudefroy, A. Gillibert, J. Guillot, F. Hammache, K. W. Kemper, E. Khan, V. Lapoux, V. Lima, L. Nalpas, A. Obertelli, E. C. Pollacco, F. Skaza, U. D. Pramanik, P. Roussel-Chomaz, D. Santonocito, J. A. Scarpaci, O. Sorlin, S. V. Stepantsov, G. M. T. Akopian, and R. Wolski, AIP Conference Proceedings **912**, 3 (2007).

[8] M. S. Golovkov, L. V. Grigorenko, A. S. Fomichev, A. V. Gorshkov, V. A. Gorshkov, S. A. Krupko, Y. T. Oganessian, A. M. Rodin, S. I. Sidorchuk, R. S. Slepnev, S. V. Stepantsov, G. M. Ter-Akopian, R. Wolski, A. A. Korsheninnikov, E. Y. Nikolskii, V. A. Kuzmin, B. G. Novatskii, D. N. Stepanov, P. Roussel-Chomaz, and W. Mittig, Phys. Rev. C **76**, 021605 (2007).

[9] H. Alfalou, Ph.D. thesis, Université de Caen, France (2007).

[10] H. Johansson, Y. Aksyutina, T. Aumann, K. Boretzky, M. Borge, A. Chatillon, L. Chulkov, D. Cortina-Gil, U. D. Pramanik, H. Emling, C. Forssn, H. Fynbo, H. Geissel, G. Ickert, B. Jonson, R. Kulessa, C. Langer, M. Lantz, T. LeBleis, K. Mahata, M. Meister, G. Mnzenberg, T. Nilsson, G. Nyman, R. Palit, S. Paschalis, W. Prokopowicz, R. Reifarth, A. Richter, K. Riisager, G. Schrieder, H. Simon, K. Smmerer, O. Tengblad, H. Weick, and M. Zhukov, Nucl. Phys. A **842**, 15 (2010).

[11] A. Matta, Ph.D. thesis, Université Paris-Sud XI (2012).

[12] N. Poppelier, L. Wood, and P. Glaudemans, Phys. Lett. B **157**, 120 (1985).

[13] E. K. Warburton and B. A. Brown, Phys. Rev. C **46**, 923 (1992).

[14] N. A. F. M. Poppelier, A. A. Wolters, and P. W. M. Glaudemans, Z. Phys. A **346**, 11 (1993).

[15] K. Hisashi and S. Hiroyuki, Nucl. Phys. A **551**, 16 (1993).

[16] T. Otsuka, R. Fujimoto, Y. Utsuno, B. A. Brown, M. Honma, and T. Mizusaki, Phys. Rev. Lett. **87**, 082502 (2001).

[17] H. Bohlen, A. Blazevic, B. Gebauer, W. V. Oertzen, S. Thummerer, R. Kalpakchieva, S. Grimes, and T. Massey, Progress in Particle and Nuclear Physics **42**, 17 (1999).

[18] A. A. Ogloblin, Z. Phys. A **351**, 355 (1995).

[19] S. Aoyama, K. Kato, and K. Ikeda, Phys. Rev. C **55**, 2379 (1997).

[20] J. Deutsch, L. Grenacs, J. Lehmann, P. Lipnik, and P. Macq, Phys. Lett. B **28**, 178 (1968).

[21] N. Vinh Mau and J. Pacheco, Nucl. Phys. A **607**, 163 (1996).

[22] Y. Kondo, T. Nakamura, Y. Satou, T. Matsumoto, N. Aoi, N. Endo, N. Fukuda, T. Gomi, Y. Hashimoto, M. Ishihara, S. Kawai, M. Kitayama, T. Kobayashi, Y. Matsuda, N. Matsui, T. Motobayashi, T. Nakabayashi, T. Okumura, H. Ong, T. Onishi, K. Ogata, H. Otsu, H. Sakurai, S. Shimoura, M. Shinohara, T. Sugimoto, S. Takeuchi, M. Tamaki, Y. Togano, and Y. Yanagisawa, Phys. Lett. B **690**, 245 (2010).

[23] M. Labiche, F. M. Marqués, O. Sorlin, and N. Vinh Mau, Phys. Rev. C **60**, 027303 (1999).

[24] G. Blanchon, N. Vinh Mau, A. Bonaccorso, M. Dupuis, and N. Pillet, Phys. Rev. C **82**, 034313 (2010).

[25] J. C. Pacheco and N. Vinh Mau, Phys. Rev. C **65**, 044004 (2002).

[26] J. Decharge and D. Gogny, Phys. Rev. C **21**, 1568 (1980).

[27] J. F. Berger, M. Girod, and D. Gogny, Comput. Phys. Commun. **63**, 365 (1991).

[28] A. Bohr and B. Mottelson, in *Nuclear Structure*, Vol. 1 (W.A. Benjamin, New York, 1969) p. 239.

[29] G. Audi, A. Wapstra, and C. Thibault, Nucl. Phys. A **729**, 337 (2003).

[30] H. Alfalou, A. Leprince, and N. Orr, Journal of Physics: Conference Series **312**, 092012 (2011).